

Periodic parabolic problems with nonlinearities indefinite in sign ^{*†}

T. Godoy [‡]- U. Kaufmann [§]
Facultad de Matemática, Astronomía y Física,
Universidad Nacional de Córdoba,
Ciudad Universitaria, 5000 Córdoba, Argentina.

December 13, 2005

Abstract

Let $\Omega \subset \mathbb{R}^N$ be a smooth bounded domain. We give sufficient conditions (which are also necessary in many cases) on two nonnegative functions a, b that are possibly discontinuous and unbounded for the existence of nonnegative solutions for semilinear Dirichlet periodic parabolic problems of the form $Lu = \lambda a(x, t) u^p - b(x, t) u^q$ in $\Omega \times \mathbb{R}$, where $0 < p, q < 1$ and $\lambda > 0$. In some cases we also show the existence of solutions u_λ in the interior of the positive cone and that u_λ can be chosen such that $\lambda \rightarrow u_\lambda$ is differentiable and increasing. A uniqueness theorem is also given in the case $p \leq q$. All results remain valid for the corresponding elliptic problems.

^{*}Partially supported by CONICET, ANPCYT, SECyT-UNC and Fundación Antorchas

[†]2000 *Mathematics Subject Classification*. 35K20, 35P05, 35B10, 35B50. *Key words and phrases*. Periodic parabolic problems, indefinite nonlinearities, elliptic problems

[‡]e-mail: godoy@mate.uncor.edu

[§]e-mail: kaufmann@mate.uncor.edu